

# Journal of Advanced Scientific Research

ISSN 0976-9595 Review Article

Available online through http://www.sciensage.info/jasr

## RANJAK PITTA AND ITS AFFILIATES IN MODERN PERSPECTIVE: A REVIEW

Rakesh Roushan\*, Pritam Moharana

Dept of Kriya Sharir, Ch. Brahm Prakash Ayurved Charak Sansthan, New Delhi, India. \*Corresponding author: rakesh3151@gmail.com

# ABSTRACT

Science of *Ayurveda* is based on *tridosha* theory. *Pitta*, one among three *doshas* plays a major role in digestion and metabolism. *Pitta dosha* is of five types namely *pachaka*, *ranjaka*, *alochaka*, *bhrajaka*, *and sadhaka pitta*. Function of *ranjaka pitta* is to provide red coloration to *rakta dhatu*. *Yakrit* (liver), *pleeha* (spleen) and *amashaya* (stomach) are the sites of *ranjaka pitta* mentioned in ancient literature. *Raktagni* and *ranjaka pitta* has an important role in the formation of *rakta dhatu*. *Sushrut* has mentioned about *sarakta meda* present inside the *alpasthi* (all bones except large bones) which is equivalent to red bone marrow and responsible for erythropoiesis. In modern physiology it is described that Liver, Spleen and Bone marrow are the site for erythropoiesis. Liver stores important factors like vitamin  $B_{12}$ , Folic acid, Iron. Red pulp of the spleen is a special reservoir of concentrated red blood cells. Intrinsic factor is a substance which is essential for absorption of vitamin  $B_{12}$  in the ileum. It is secreted by the parietal cell of stomach. Erythropoiesis. In this article we are intended to identify the entities of modern medical sciences that have resemblance with *Ranjaka pitta* on the basis of its physiological functions. For this study, the basic materials have been collected from the *Ayurvedic* classics with the available commentaries, as well as text books of modern medical science for better understanding of the concept and its comparison with contemporary science.

Keywords: Erythropoiesis, Rakta dhatu, Raktagni, Ranjak pitta

## 1. INTRODUCTION

Many Ayurvedic principles have been described in Ayurvedic Samhitas. Among them theory of tridoshas is an important principle. *Tridosha* is responsible for homeostasis of human body. Vata, pitta and kapha doshas is not represented by a single entity in our body. We can understand these concepts on the basis of its functional aspects [1]. Again the function of each *dosha* is unique. All three doshas are equally important. A single dosha executes many actions. Therefore we cannot represent a single structure for the specific function of each *dosha*. All these are equally important and do their functions independently or sometimes with co-operation of other. Pitta dosha one among three dosha is responsible for digestion and metabolism in human body. Wherever any type of chemical reaction takes place in the body there should be involvement of enzyme, hormone, or neurotransmitter as a result there is production of heat or ATP like energy producing substance. Those substances may be considered as pitta [2].

*Pitta dosha* is present in cellular level, single system level, and organization level. *Pitta dosha* do not move in the body independently. It is circulated all over the body with the help of *vata dosha*. In modern physiology it has been mentioned that the basic theory of body's control system allow the functional system to operate in support of one another [3]. The homeostatic mechanism involves Nervous system, Hormone system and Immune system. Similarly all these three *doshas* have equal importance and they together maintain homeostasis in our body. Here the function of *pitta dosha* may represent the enzymes, endocrine or neuroendocrine system in organization level.

*Pitta dosha* are of five subtypes on the basis of location namely *paachak*, *ranjak*, *saadhak*, *alochaka*, *bhrajaka pitta*. One of the most important liquid (*drava*) *dhaatu* in our body is called *rakta dhaatu*. It is essential for nourishment and equilibrium. *Raktagni* (*dhatwagni*) and *ranjak pitta* helps simultaneously in the formation of *rakta dhatu*. There is a need of proper understanding for the formation of *rakta dhatu* in modern perspective. *Ranjak pitta* and *raktagni* cannot be represented by a single entity at all the time because *Ayurveda* is the science based on the concept of functional understanding.

There is no specific correlation of *Ranjaka pitta* mentioned in *Ayurvedic* literature in terms of supporting modern literature. It seems to be a problem found in student life, particularly first year of Bachelor of *Ayurvedic* Medicine and Surgery to understand about the concept of *Ranjaka pitta*. Increased demand of *Ayurveda* science in

the present scenario is required to understand the depth of *Ayurvedic* principles on criterion of modern medical science in an easy mode. In this review we are trying to identify anatomical structures based on its physiological functions retrospectively described under the function of *Ranjaka pitta*.

Site and	l func	tion of	ran	jaka	pitta	by	different	achar	ya:
									/

	Sushruta Samhita [4]	Astanga Hridaya [5]	Astanga Samgraha [6]	Sharngadhar Samhita [7]
Sthana (Location)	Yakrit (liver), pliha (spleen)	Aamashaya (stomach)	Aamashaya (stomach)	<i>Yakrit</i> (Liver)
Karma (Function)	To impart colour to rasa	To impart colour of	To impart colour of	Converts rasa into rakta
	dhatu	rasa dhatu	rasa dhatu	

#### 2. RANJAKA PITTA AND RAKTAGNI

Aahar rasa is final absorbed product of food digested by jatharagni. From aahar rasa subsequent dhatus are formed and nourished by three laws [8]. Commentator provides three laws for the formation and nourishment of *dhatus*. Those are Kshiradadhi nyaya, Khalekapota nyaya, Kedarikulya nyaya. Three laws work together simultaneously for the replenishment of dhatus. Kshiradadhi nyaya work for the formation of dhaatu where as kedarikulya nyaya and khalekapota nyaya work for the nourishment of dhaatu. Jatharagni act upon aahar and is converted into aahar rasa and mala [9]. In kshiradadhi nyaya Acharya Charak has mentioned rasagni act upon rasa dhatu and is converted into rakta dhatu [10] whereas in Sushrut samhita Raktagni act upon anu rakta which is formed from rasa dhatu by the action of rasagni and help in the formation of rakta dhatu [11]. In kedarikulya nyaya and khalekapota nyaya rakta sadharmi ansa (nutrient specific to dhatu) of aahra ras (nutrient product) nourishes the formed *dhatus*. Blood should be regarded as pure when its color resembles red-gold, indragopa (scarlet fly), red lotus, mahavar, and gunja fruit (abrus precatorius) [12]. The function of *raktagni* is to synthesize the colorless *rakta* dhaatu from anu part of rasa dhaatu.

Again Acharya Charak has mentioned that the teja portion of rasa is mixed with ushma of pitta (ranjak pitta) that is helpful for the coloration of rakta dhatu [13]. Acharya Dalhana in the commentary of Sushrut Samhita has mentioned tejarupa ranjak pitta is mixed with avyapanna vikar rahit rasa (diseased free rasa dhatu) to form rakta dhatu [14]. Acharya sushrut has named all subtype of pitta by the name of agni. Acharya sushruta has mentioned the location of Ranjaka pitta as yakrit (liver) and pliha (spleen). It helps in the formation of *rakta dhatu*. *Vagbhatta* has mentioned its location as *amashaya* (stomach).According to all these views *rasa dhatu* is contributed in the formation of *rakta dhatu* with the help of *ranjaka pitta*. The function of *ranjaka pitta* is to color the *rasa dhatu* to form *rakta dhatu*. So, both *ranjaka pitta* and *raktagni* has an important role in the formation of *rakta dhatu*.

The root of raktavaha srotas is yakrit (liver) and pliha (spleen) and raktavahi dhamani (blood vessels). Acharya Charak has described srotas are the channels which transport their respective *dhatus* which means *yakrit*, *pliha* and raktavahi dhamani carries the rakta dhatu to hridaya through saman vayu [15]. Rakta dhatu is formed elsewhere in the body. The third kala is called as medodhara kala. Medas is found in the abdomen and small bones of all living beings. The long bone contains bone marrow (instead of medas). Further, Acharya Sushruta has classified bones as sthula asthi and other then sthoolasthi. Acharya Dalhan has commented alpa asthi for bones other then sthoolasthi. Bone marrow is found especially (visheshtah) inside the sthula asthi (large bones) and sarakta medas (red bone marrow) is found in bones other then sthoolasthi (alpa asthi) [16].

Bone marrow is specifically present in *sthulasthi*. As per *arthapatti tantrayukti* we can say that some less amount of other type of *majja* may be present and this other type of marrow may be red bone marrow. In some large bones like humerus and tibiae red bone marrow is present in the proximal portion. *Acharya Sushrut* has explained red bone marrow is present in the proximal portion. *Acharya Sushrut* has explained red bone marrow is present in *alpasthi* (bones like vertebrae, sternum, ribs, ilea).

## 3. MODERN ASPECTS

Erythropoiesis is the process of formation of red blood cells. In the early week of embryonic life red blood cells are produced in yolk sac. During the middle trimester liver is the main organ for production of red blood cells. Spleen and lymph node is also responsible for the production of red blood cell during this period. During the last month of gestation and after birth red blood cells are produced in bone marrow. Bone marrow of all bones produces red blood cells until 5 years old. The bone marrow of long bones except the proximal portion of humeri and tibiae becomes fatty and produces no more red blood cells after 20 years. After this age most of red cells are produced in marrow of membranous bone like vertebrae, sternum, ribs and ilia. The marrow of these bones becomes less productive as age increases [17].

All the cells of circulation blood are derived from pluripotential hematopoietic stem cell in bone marrow. Growth and reproduction of different stems cells are controlled by multiple proteins called growth inducers. Four major growth inducers have been described. Interleukin-3 promotes growth and reproduction of all different type of committed stem cells whereas others induce growth of only specific type of cell. Differentiation inducers are the proteins that promote differentiation of cells. Formation of growth inducers and differentiation inducers is controlled by factors outside the bone marrow [18].

Pluripotent hematopoietic stem cell is converted to committed stem cell. The different committed stem cell grows to produce specific type of blood cells. A committed stem cell called colony forming uniterythrocyte (CFU-E) produces erythrocytes. Under appropriate stimulation CFU-E stem cells is converted to proerythroblast. Once proerythroblast is formed it divides into multiple times to form many mature red blood cells. Proerythroblast is converted into basophil erythroblast/early normoblast. At this time the cell is accumulated with very little hemoglobin. In the next step early normoblast is converted into polychromatophil erythroblast/intermediate normoblast then to orthochromatic erythroblast then to reticulocyte. In this step the cells become filled with haemoglobin to a concentration of 34%. The nucleus condenses to small size and its final remnant is absorbed or extruded from the cell. During this reticulocyte stage, the cells pass from the bone marrow into the blood capillaries by diapedesis. The cell is then called mature erythrocyte [19].

#### 4. SPECIFIC BLOOD RESERVOIR

Spleen can release 100 ml of blood into other area of circulation by decreasing its size sufficiently. The sinuses of liver can release about 450 milliliters of blood into remainder of circulation. Large abdominal veins can contribute 300 ml; venous plexus can contribute several hundred milliliters. Spleen has two areas for storing blood venous sinuses and pulp. The red pulp of the spleen is a special reservoir that contains large quantities of concentrated red blood cells. These red blood cells can be expelled into general circulation whenever sympathetic nervous system becomes excited and causes the spleen and its vessels to contract. 50 ml of concentrated red blood cell can be released into circulation, raising the hematocrit 1-2 % [20].

## 5. SITES OF RANJKA PITTA

The site of ranjak pitta is yakrit (Liver), pliha (Spleen), amashaya (Stomach) which is well proved by our Ayurveda masters. In modern literature it is mentioned that blood is formed in liver and spleen during intrauterine life. Hepatic stem cells in fetal liver migrate to bone marrow and become the major site for production of blood after birth. Liver is store house for Vitamin B<sub>12</sub>. Folic acid and iron. It is needed for the maturation of erythrocyte. The red pulp of the spleen is a special reservoir that contains large quantities of concentrated red blood cells. In liver and spleen diseases like hepatomegaly and splenomegaly there is reduction of hemoglobin concentration which denotes that there is importance of liver and spleen for the formation of blood. It is also seen that when liver and spleen are in normal condition still there is reduction in hemoglobin. It is due to the absence of intrinsic factor which is secreted from parietal cells of stomach (amashaya) and is required for the absorption of vitamin  $B_{12}$ . In ancient literature it is also mentioned that when there excessive blood loss, one should drink blood with honey or he should eat goat's fresh liver with its contained bile [21]. So these three structures yakrit (liver), pleeha (spleen), and amashaya (stomach) help in maturation of erythrocyte and proliferation of erythroblastic cells which supports our Ancient Acharya. In modern science it is described after 20 years of age

blood formation occurs in red bone marrow of vertebrae, sternum, ilea, ribs and proximal portion of humerus and tibbie. The actual site of *rakta dhatu* formation (erythropoiesis) is not specified. *Acharya Sushruta* has classified bones as *sthula asthi* and other then *sthoolasthi*. *Acharya Dalhan* has commented *alpa asthi* for bones other then *sthoolasthi*. Bone marrow is found especially (visheshtah) inside the sthula asthi (large bones) and sarakta medas (red bone marrow) is found in bones other then sthoolasthi (alpa asthi) [22].

Bone marrow is present in *sthulasthi* in significant quantity. As per *arthapatti tantrayukti* we can say that some less amount of other type of *majja* may be present and this other type of marrow may be red bone marrow. In some large bones like humerus and tibiae red bone marrow is present in the proximal portion. *Acharya Sushrut* has explained red bone marrow is present in the proximal portion. *Acharya Sushrut* has explained red bone marrow is present in *alpasthi* (bones like vertebrae, sternum, ribs, ilea). *Rakta sadharmi ansa* is present in *aahar* which is required for the nutrition of *rakta dhatu*. Food content which is required to nourish and proliferation the pluripotent stem cells may be called as *rakta sadharmi ansa*.

# Correlation of functions of raktagni and ranjaka pitta:

<i>Raktagni</i> and its	Growth inducers, Differentiation			
correlation	inducers, Erythropoietin,			
	Norepinephrine, Epinephrine,			
	Prostaglandin, Androgen			
	hormones			
Ranjak pitta and its	Intrinsic factor, Vitamin B <sub>12</sub> ,Folic			
correlation	acid, Iron, Copper			

# 6. GROWTH INDUCERS AND DIFFERENTIATION INDUCERS

The cells of circulation blood are derived from pluripotential hematopoietic stem cell in bone marrow. Growth and reproduction of different stems cells are controlled by multiple proteins called growth inducers. Differentiation inducers are the proteins that promote differentiation of cells. As the function of *raktagni* is to synthesize the *rakta dhatu* so the function of Growth inducers and Differentiation inducers comes under the function of *Raktagni*.

## 6.1. Erythropoietin

Conditions that decrease the partial pressure of oxygen in the blood increase the rate of red blood cell production. It helps in formation of growth inducer and differentiation inducer as a result there is increased number of erythrocytes. This process is carried out by erythropoietin. So Erythropoietin enhances red blood cell production. Thus the function of Erythropoietin comes under the function of *Raktagni*.

# 6.2. Norepinephrine, Epinephrine, Prtostaglan din, Androgen

Norepinephrine, epinephrine and several of prostaglandin stimulate kidney for erythropoietin production. The effect of testosterone to increase red blood cell production may be at least partly indirect because of the increased metabolic rate that occurs after testosterone administration. So norpeinephrine, epinephrine, prostaglandin, testosterone have an important role in production. erythrocyte So the function of norepinephrine, epinephrine, prtostaglandin, and androgens are come under the functions of raktagni.

# 6.3. Intrinsic Factor

The parietal cells of the gastric gland secrete a glycoprotein called intrinsic factor, which combine with vitamin  $B_{12}$  in food and makes it available for absorption by the gut. Intrinsic factor binds tightly with vitamin  $B_{12}$ . In this bound state vitamin  $B_{12}$  is protected from digestion by gastrointestinal secretion. Intrinsic factor binds to specific receptor site on the brush border membrane of the mucosal cells of ileum. A common cause of red blood cell maturation failure is failure to absorb vitamin  $B_{12}$  from gastrointestinal tract. In diseases like atrophic gastric mucosa stomach fails to produce normal gastric secretions. Lack of intrinsic factor decreases the availability of vitamin  $B_{12}$  and failure in maturation of RBCs. So, the functions of intrinsic factor come under the function of *ranjak pitta*.

# 6.4. Vitamin B<sub>12</sub>, Folic Acid

Vitamin B<sub>12</sub> and folic acid plays an important role for final maturation of red blood cell. These two are essential for the synthesis of DNA because this is required for the formation of thymidine triphosphate which is the essential building blocks of DNA. Lack of either vitamin  $\boldsymbol{B}_{12}$  or folic acid causes abnormal and diminished DNA consequently failure of nuclear maturation and cell division. The erythroblastic cells of the bone marrow due to failing of proliferation rapidly produce larger than normal red cell called macrocyte. These poorly formed cells are capable of carrying normal oxygen but their lifespan will have shorter one third to half. Vitamin  $B_{12}$  is transported into the blood during next few hours by the process of pinocytosis, carrying intrinsic factor and vitamin  $B_{12}$  together through the membrane. Once vitamin  $B_{12}$  is absorbed from GIT it is first stored in liver and then released slowly as needed by the bone marrow. Folic acid is normal constituent of green vegetable, fruits, and meats (especially liver). Failure of maturation of

RBC is also caused by the deficiency of folic acid. In certain case like sprue there is difficulty of absorption of folic acid and vitamin  $B_{12}$  which causes failure of maturation of RBC. Therefore function of vitamin  $B_{12}$ , and folic acid come under the function of *ranjak pitta*.

## 6.5. Iron

Iron has an important role in formation of hemoglobin. When iron is absorbed from the small intestine with the help of vitamin C it immediately combines in the blood plasma with a beta globulin (apotransferrin) to form transferrin which is then transported in the plasma. In the cell cytoplasm iron combines with apoferritin to form ferritin. The hepatic cells contain large amounts of a protein called *apoferritin*, which is capable of combining reversibly with iron. Therefore, when iron is available in the body fluids in extra quantities, it combines with apoferritin to form ferritin and is stored in this form in the hepatic cells until needed elsewhere. This iron stored as ferritin is called storage iron. When quantity of iron in the plasma fall low, iron in the ferritin storage pool is removed easily and transported in the form of transferrin in the plasma to the areas of the body where it is needed. Transferrin molecule binds strongly with receptors in the cell membrane of erythroblast. There the transferrin delivers the iron directly to the mitochondria where heme is synthesized. People who do not have adequate quantities of transferrin in their blood can cause severe hypochromic anemia. So iron has an important role in maturation of RBCs. That's why functions of iron come under the function of *ranjak pitta*.

#### 6.6. Copper

Copper profoundly influences the absorption of iron. This suggests that a copper protein may be involved in the mechanism by which iron is absorbed from the gastrointestinal tract. The daily turnover rate of iron through the plasma and into the red cells is increased rather than decreased in copper-deficient swine suggests that movement of iron within the body is not reduced. The relation of copper to iron metabolism must be in some way concerned with the role of copper in red cell synthesis. Anemia develops in the absence of copper because of the limited capacity of the marrow to produce cells. It is suggested that deficiency of the erythrocyte copper compound erythrocuprein, is related to these alterations in erythrocyte production and survival [23]. Thus function of copper come under the function of ranjaka pitta.

#### 7. DISCUSSION

Basically Vata, Pitta, Kapha constitute three regulatory systems i.e. nervous, endocrine and immune system respectively of all living system. Among five types of pitta, Ranjaka pitta has an important role for the coloration of rakta dhatu. Raktagni (one type of Dhatavagni) helps in formation of rakta dhatu. Raktagni acts on anu rakta (pluripotent stem cells) and convert it into rakta dhatu (blood cells). Ranjak pitta during the process of hemopoiesis provides coloration (hemoglobin) to rakta dhatu.

Acharya Sushrut has mentioned the site of ranjaka pitta as yakrit and pliha, Acharya Sharngadhar has metioned it as yakrit and Acharya vagbhatta has mentioned as amashyaya (Stomach). Acharya Charak has not mentioned the site for ranjak pitta. Acharya Sushrut and Vagbhatta have mentioned the function of ranjaka pitta is to impart coloration whereas Acharya Sharngadhar has mentioned the function of ranjaka pitta is to convert rasa dhatu into rakta dhatu. Acharya Charak has mentioned teja portion of ahara rasa is mixed with ushma of pitta (ranjak pitta) that is helpful for the coloration of rakta dhatu.

In modern physiology hormone like erythropoietin, androgen, thyroid hormone, corticosteroid hormone vitamins like vitamin B<sub>12</sub>, folic acid, pyridoxine, vitamin C (helps iron absorption), minerals like iron and copper, norepinephrine, epinephrine, intrinsic factor, Growth inducers, differentiation inducers are the major factors that is responsible for erythropoiesis and maturation of RBC. From the above discussion functions of Growth inducers, differentiation inducers, erythropoietin, epinephrine, norepinephrine, prostaglandin and testosterone may be correlated with the functions of raktagni. Functions of Intrinsic factor, Vitamin B<sub>12</sub>, folic acid, iron and copper may be correlated with the functions of ranjak pitta.

#### 8. INADEQUECIES

Across all Ayurvedic textbooks, the following inadequacies are found in relation to blood physiology.

- There is no direct reference for the site of production of *rakta dhatu* (blood) in fetal life as well as adult life which are clearly described in modern literature.
- Name of the specific bones is not mentioned where *sarakta meda* (red bone marrow) is present.
- Acharya Sushrut has mentioned sarkta meda are present in asthis (alpasthi) other than sthulasthi. Examples of sthulasthi and alpasthi are not mentioned in the text.
- Function of *sarkta meda* is not specified.

- Acharya charak has mentioned the role of rasagni in the production of rakta dhatu where as Acharya sushrut has mentioned the role of raktagni in the production of rakta dhatu.
- Stage wise haemopoiesis is not mentioned clearly by our ancient *acharya*.
- Location of Raktagni (dhatavagni) is not mentioned.
- In most of the text books the word *prasadaja* is considered for *dhatu nirman*. *Dhatu poshana* is the actual meaning for *prasadaja* of *dhatu* [24].

## 9. CONCLUSION

*Pitta dosha* is divided into five types namely *pachaka*, *ranjaka*, *sadhaka*, *alochaka*, *and bhrajaka*. The site of *ranjak pitta* as described in ancient literature is *yakrit*, *pliha*, *and amashaya*. *Ayurveda* is based on functional understanding. From the above discussion functions of *Raktagni* may be correlated with Growth and differentiation inducers, erythropoietin, norepinephrine, epinephrine, prostaglandin, and testosterone. Functions of *Ranjak pitta* may be correlated with intrinsic factor, vitamin B<sub>12</sub>, Folic acid, Iron and copper. There is a need of further research to evaluate in detail of all other *doshas*. So, that student joining BAMS can understand easily the basic concepts of *rakta dhatu* and *ranjaka pitta*.

## **10. REFERENCES**

- Moharana P, Rawat N, Roushan R. International Journal of basic and applied research, 2018; 8(9):897-903.
- Moharana P, Roushan R. Int. J. Res. Ayurveda Pharm., 2019; 10(1):18-20.
- Hall JE, Guyton AC. General Physiology. In: Vaz M, editors. Textbook of medical physiology. 2<sup>nd</sup> ed. New Delhi (India): Elseveir; 2018. p. 2-9.
- Sushrut. Vranaprashna Adhyaya. In: Shastri AD, editors. Susruta Samhita. Revised edition. Varanasi (India): Chaukhamba Sanskrit Sansthan; 2016. p. 112-122.
- Vagbhatta. Doshabhediya Adhyaya. In: Tripathy BN, editors. Astanga Hridaya. Revised edition. Delhi (India): Chaukambha Sanskrit Pratisthan; 2012. p. 170-184.
- Vaghbhatta. Doshabhediya Adhyaya. In: Gupta AD, editors. Astanga Sangraha. Revised edition. Varanasi (India): Chaukambha Krishnadas Academy; 2016. p. 159-166.
- 7. Sharngadhar. Aaharadi gati adhyaya. In: Shastri DD, editors. Shrangadhar Samhita. Revised edition.

Varanasi (India): Chukhamba Surbharati Prakashan; 2002. p. 83-94.

- Moharana P, Roushan R. Int. J. Ayu. Pharm. Chem., 2018; 9(2):537-547.
- Rawat N, Roushan R. Int. J. Res. Ayurveda Pharm., 2019; 10(1):28-32.
- Chakrapani. Grahani Chikitsa adhyaya. In: Gaur BL, editors. Ayurveda deepika on Charak Samhita. 1<sup>st</sup> ed. New Delhi (India): Rastriya ayurveda vidyapeeth; 2014. p. 796-875.
- Dalhan. Sonita varnaniya adhyaya. In: Prasad BB, editors. Nibandha Sangraha on Sushrut Samhita. Revised edition. New Delhi (India): Rastriya ayurveda vidyapeeth; 2002. p.133-146.
- Charak, Vidhisonita adhyaya. In: Pandey K, Chaturvedi G, editors. Charaka Samhita. Revised edition. Varanasi (India): Chaukambha Bharati Academy; 2015. p. 442-456.
- Charak, Vidhisonita adhyaya. In: Pandey K, Chaturvedi G,editors. Charaka Samhita. Revised edition. Varanasi (India): Chaukambha Bharati Academy; 2015. p. 442-456
- Sushrut, Vranaprashna Adhyaya. In: Shastri AD, editors. Susruta Samhita. Revised edition. Varanasi (India): Chaukhamba Sanskrit Sansthan; 2016. p. 112-122.
- Moharana P, Roushan R. Int. J. Ayu. Pharm. Chem., 2018; 9(2): 188-197.
- Sushrut. Garbhavyakarana shareera. In: Shastri AD, editors. Susruta Samhita. Revised edition. Varanasi (India): Chaukhamba Sanskrit Sansthan; 2016. p. 36-54.
- Hall JE, Guyton AC. Blood and its constituents. In: Kurpad A, editors. Textbook of medical physiology. 2<sup>nd</sup> ed. New Delhi (India): Elseveir; 2018. p. 109-167.
- Hall JE, Guyton AC. Blood and its constituents. In: Kurpad A, editors. Textbook of medical physiology. 2<sup>nd</sup> ed. New Delhi (India): Elseveir; 2018. p. 109-167.
- Hall JE, Guyton AC. Blood and its constituents. In: Kurpad A, editors. Textbook of medical physiology. 2<sup>nd</sup> ed. New Delhi (India): Elseveir; 2018. p. 109-167.
- Hall JE, Guyton AC. Cardiovascular physiology. In: Vaz M, editors. Textbook of medical physiology. 2<sup>nd</sup> ed. New Delhi (India): Elseveir; 2018. p. 171-323.
- 21. Charak. Arsha Chikitsa Adhyaya. In: Pandey K, Chaturvedi G, editors. Charaka Samhita. Revised

edition. Varanasi (India): Chaukambha Bharati Academy; 2015. p. 418-452

- Sushrut, Garbhavyakarana shareera. In: Shastri AD, editors. Susruta Samhita. Revised edition. Varanasi (India): Chaukhamba Sanskrit Sansthan; 2016. p. 36-54.
- 23. Cartwright GE, Gubler CJ. National Academy of Sciences-National Research Council, 1958; 11(3):100.
- Chakrapani. Grahani Chikitsa adhyaya. In: Gaur BL, editors. Ayurveda deepika on Charak Samhita. 1<sup>st</sup> ed. New Delhi (India): Rastriya ayurveda vidyapeeth; 2014. p. 796-875.